



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: September 9, 2014

ACCEPTED: November 11, 2014

PUBLISHED: November 27, 2014

Order α' heterotic domain walls with warped nearly Kähler geometry

Alexander S. Haupt,^{a,b} Olaf Lechtenfeld^a and Edvard T. Musaev^{a,c,d}

^a*Institut für Theoretische Physik and Riemann Center for Geometry and Physics,
Leibniz Universität Hannover, Appelstraße 2, 30167 Hannover, Germany*

^b*Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut),
Am Mühlenberg 1, 14476 Potsdam, Germany*

^c*Université de Lyon, Laboratoire de Physique, UMR 5672, CNRS,
École Normale Supérieure de Lyon, 46 allée d'Italie, 69364 Lyon CEDEX 07, France*

^d*National Research University Higher School of Economics, Faculty of Mathematics,
20, Myasnitskaya street, 101000 Moscow, Russia*

E-mail: alexander.haupt@itp.uni-hannover.de,
olaf.lechtenfeld@itp.uni-hannover.de, emusaev@hse.ru

ABSTRACT: We consider (1+3)-dimensional domain wall solutions of heterotic supergravity on a six-dimensional warped nearly Kähler manifold X_6 in the presence of gravitational and gauge instantons of tanh-kink type as constructed in [1]. We include first order α' corrections to the heterotic supergravity action, which imply a non-trivial Yang-Mills sector and Bianchi identity. We present a variety of solutions, depending on the choice of instantons, for the special case in which the $SU(3)$ structure on X_6 satisfies $W_1^- = 0$. The solutions preserve two real supercharges, which corresponds to $\mathcal{N}=1/2$ supersymmetry from the four-dimensional point of view. Besides serving as a useful framework for collecting existing solutions, the formulation in terms of dynamic $SU(3)$ structures utilized here allows us to obtain new solutions in as yet unexplored corners of the instanton configuration space. Our approach thus offers a unified description of the embedding of tanh-kink-type instantons into half-BPS solutions of heterotic supergravity where the internal six-dimensional manifold has a warped nearly Kähler geometry.

KEYWORDS: Flux compactifications, Solitons Monopoles and Instantons, Supergravity Models

ARXIV EPRINT: [1409.0548](https://arxiv.org/abs/1409.0548)

Contents

1	Introduction	1
2	Geometry of the domain-wall background	3
3	Yang-Mills sector	8
3.1	Yang-Mills instantons on $\mathbb{R} \times X_6$	8
3.2	Bianchi identity	10
4	Warped nearly Kähler domain wall	11
4.1	Static versus dynamic $SU(3)$ structures on X_6	11
4.2	Embedding of the instanton in the ten-dimensional solution	12
4.3	Equations of motion	13
4.4	Summary of the system of equations	14
5	Explicit solutions with $W_1^- = 0$	15
6	Conclusions and outlook	23
A	Conventions	25

1 Introduction

The problem of finding solutions of ten-dimensional supergravities with a compact manifold filling internal directions is of manifest importance for phenomenological applications (for reviews on that subject, see, for example, [2–6]). Most of the variety of different vacua in string theory comes from the choice of the internal manifold, the simplest example of which is a flat torus. However, toroidal compactifications lead to lower-dimensional theories that are, in some sense, “too simple” in that they typically do not entail realistic phenomenology. In particular, toroidal compactifications of minimally supersymmetric ten-dimensional theories to four dimensions yield $\mathcal{N} = 4$ theories that are non-chiral [7]. Less trivial examples of manifolds that lead to interesting lower-dimensional physics are given by manifolds with special geometry such as Calabi-Yau (or, more generally, $SU(3)$ structure) manifolds [8]. An important feature of these manifolds is that they preserve less supersymmetry, thus leading to more realistic models. This is a direct consequence of the holonomy principle, which states that the parallel spinor equation

$$\nabla \epsilon = 0 \tag{1.1}$$

has m solutions ϵ if and only if the holonomy group of ∇ is contained in the joint stabilizer subgroup of m spinors, which in turn is related to the G structure of the manifold. Then, m defines the amount of supersymmetry preserved by the background.

In this paper, we study heterotic supergravity, that is the low-energy limit of heterotic string theory, which was first constructed in [9–11]. Heterotic supergravity consists of $\mathcal{N} = 1$, $D = 10$ supergravity coupled to super Yang-Mills theory. The ingredients are a ten-dimensional manifold \mathcal{M} , equipped with a Lorentzian metric \hat{g} , an NS 3-form \hat{H} , a dilaton $\hat{\phi}$ and a gauge connection ${}^A\hat{\nabla}$, with gauge group $\text{SO}(32)$ or $E_8 \times E_8$. It was shown in [9] that the anomaly cancellation condition of ten-dimensional super Yang-Mills theory coupled to $\mathcal{N} = 1$, $D = 10$ supergravity can be written as a Bianchi identity on \hat{H} ,

$$\hat{d}\hat{H} = \frac{\alpha'}{4} \text{Tr}(\hat{F} \wedge \hat{F} - \tilde{R} \wedge \tilde{R}) , \quad (1.2)$$

where \hat{d} is the ten-dimensional exterior derivative, \hat{F} is the curvature 2-form of the gauge connection ${}^A\hat{\nabla}$ and \tilde{R} is the curvature 2-form of a connection $\tilde{\nabla}$. Different connections used in the anomaly cancellation condition (1.2) correspond to different renormalization schemes [12] and there is some discussion about the correct choice of the curvature \tilde{R} in the literature (see, for example, [13] and references therein). In particular, string theory appears to prefer the choice $\tilde{\nabla} = {}^+\hat{\nabla}$ [14, 15], with $\pm\hat{\nabla}$ being the metric compatible connections on the tangent bundle of \mathcal{M} with torsion $\pm\hat{H}$. From a purely supergravity point of view, the connection $\tilde{\nabla}$ is determined by imposing the instanton equation $\tilde{R} \cdot \epsilon = 0$ [13]. For the purpose of this paper, we will adopt the latter point of view.

In the case of vanishing NS 3-form flux, $\hat{H} = 0$, the internal manifold should be Ricci-flat and Kähler. Such a solution typically does not stabilize all Kähler moduli. Owing to the different scale properties of the terms on the opposite sides of the equation above, a solution with non-zero NS 3-form flux breaks scale invariance and is thus capable of stabilizing the Kähler moduli [16]. In the present paper, we construct order α' solutions with non-zero NS 3-form flux that preserve $\mathcal{N} = 1/2$ supersymmetry (that is two real supercharges) in 1+3 external dimensions. The usual $\mathcal{N} = 1$ supersymmetry (that is four real supercharges) implied by the BPS equations is halved by the presence of a domain wall.

At the zeroth order in α' , the BPS equations are solved by $\mathcal{M} = \mathbb{R}^{1,2} \times c(X_6)$ with vanishing NS 3-form flux $\hat{H} = 0$ and $\hat{\phi} = \text{const}$. Here, $c(X_6)$ is the metric cone over a six-dimensional nearly Kähler manifold X_6 . At the first order in α' , the BPS equations can be solved by choosing the gauge connection to be ${}^A\hat{\nabla} = {}^{\text{LC}}\nabla$ and $\hat{H} = 0$, $\hat{\phi} = \text{const}$ [1], where ${}^{\text{LC}}\nabla$ is the Levi-Civita connection on $c(X_6)$. Less trivial solutions with $\hat{H} \neq 0$ can be obtained if the gauge field is chosen to be an instanton [1, 17–23]. In our analysis, we exploit the instanton solution of [1] and reformulate it in the framework of dynamic $\text{SU}(3)$ structures. Moreover, we consider previously unexplored combinations of instanton configurations thereby extending results of [1, 22, 23]. In addition, we reproduce some of the solutions found in [1, 22, 23], in special corners of our instanton configuration space. Finally, in the $\alpha' \rightarrow 0$ limit, our construction becomes a sub-sector of the more general zeroth order analysis of [24].

The paper is organized as follows. In the next section, we review the G structure formalism for solving the BPS equations of heterotic supergravity, as developed in [24, 25]. We also introduce our ansatz for the metric, the NS 3-form and the dilaton and discuss the zeroth order in α' solution. At the first order in α' , the gauge field \hat{F} couples to the other fields and therefore becomes non-trivial. The subject of section 3 is to review the construction of a certain type of seven-dimensional Yang-Mills instantons that were found in [1]. These instantons are employed in order to solve the Yang-Mills sector of the theory. In section 4, we combine the seven-dimensional Yang-Mills instantons with the other fields in order to lift the zeroth order solution of section 2 to a fully 10-dimensional solution that is valid at the first order in α' . This involves solving the Bianchi identity and a careful treatment of the equations of motion up to this order. We will also unveil a subtle relationship between static and dynamic $SU(3)$ structures on the six-dimensional compact part X_6 of the ten-dimensional space-time manifold, which is a consequence of the warping included in the metric ansatz. In section 5, we present explicit solutions assuming that $W_1^- = 0$, that is the torsion class W_1 of the dynamic $SU(3)$ structure on X_6 has vanishing imaginary part. The precise dynamics of the solution depends on the choice of instanton configurations. Our solutions include, in a unified description, special cases of [1, 22, 23] and some new ones. We end the main body of the paper by providing a few concluding remarks in section 6. Finally, our conventions for indices and normalizations are summarized in appendix A.

2 Geometry of the domain-wall background

For a background with vanishing fermionic vacuum expectation values to be supersymmetric, the supersymmetry transformations of the corresponding fermionic fields must vanish. This implies certain conditions on the background, known as BPS equations. For the fermionic content of heterotic supergravity, one finds that the BPS equations up to and including terms of order α' are given by

$$\begin{aligned} -\hat{\nabla}\epsilon &= 0, \\ \left(\hat{d}\hat{\phi} - \frac{1}{2}\hat{H}\right) \cdot \epsilon &= 0, \\ \hat{F} \cdot \epsilon &= 0, \end{aligned} \tag{2.1}$$

for a Majorana-Weyl spinor ϵ . Here and in the following, hatted objects denote ten-dimensional quantities. The conventions used in this paper are summarized in appendix A.

We are interested in the background given by a four-dimensional domain wall with six internal directions filled by a compact manifold X_6 with $SU(3)$ structure. We *a priori* choose the following metric ansatz,

$$\hat{g} = e^{2A(x^m)} \left(\eta_{\alpha\beta} dx^\alpha dx^\beta + e^{2\Delta(x^u)} dx^3 dx^3 + g_{uv}(x^m) dx^u dx^v \right). \tag{2.2}$$

The world-volume of the domain wall is parametrized by the coordinates x^α with $\alpha \in \{0, 1, 2\}$. The coordinate x^3 is chosen to be transverse to the domain wall and will henceforth also be denoted by y . The orthonormal frame on the internal six-dimensional manifold

X_6 is given by $\{e^{\underline{u}}\} = \{e^{\underline{u}}_u dx^u\}$ with $u \in \{4, 5, \dots, 9\}$ and underlined indices denote tangent space (local Lorentz) indices. Finally, the set of coordinates $\{x^m\} = \{x^3, x^u\}$ combines all of the directions transverse to the domain wall world-volume. To summarize, the total ten-dimensional space-time manifold \mathcal{M} locally splits as

$$\mathcal{M} = \mathbb{R}^{1,2} \times \mathbb{R} \times X_6, \quad (2.3)$$

with a flat metric $\eta_{\alpha\beta} = \text{diag}(-1, 1, 1)$ on $\mathbb{R}^{1,2}$ and a general metric $g_{uv}(x^m)$ compatible with a y -dependent $\text{SU}(3)$ structure on X_6 .

As a starting point of our further analysis, we shall briefly repeat here the derivation of G structures consistent with the above ansatz, following [24, 25]. First, the Killing spinor ϵ is decomposed according to our metric ansatz as

$$\epsilon(x^\alpha, x^m) = \rho(x^\alpha) \otimes \eta(x^m) \otimes \theta, \quad (2.4)$$

where ρ is the covariantly constant spinor on the world-volume $\mathbb{R}^{1,2}$ of the domain wall, θ is an eigenvector of the third Pauli matrix and η is a covariantly constant Majorana spinor on the seven-dimensional space $X_7 := \mathbb{R} \times X_6$. The spinor ρ has two real components, which corresponds to the two real supercharges that our background preserves. In four-dimensional terminology this corresponds to $\mathcal{N} = 1/2$ supersymmetry.

We would like to preserve (1+2)-dimensional Lorentz invariance on the domain wall world-volume. This restricts $\hat{\phi}$ and \hat{H} such that

$$\partial_\alpha \hat{\phi} = 0, \quad \hat{H}_{\alpha mn} = 0, \quad \hat{H}_{\alpha\beta n} = 0. \quad (2.5)$$

Hence, the only non-zero components of the NS 3-form flux are \hat{H}_{yuv} , \hat{H}_{uvw} and $\hat{H}_{\alpha\beta\gamma} = \ell \epsilon_{\alpha\beta\gamma}$ with $\epsilon_{\alpha\beta\gamma}$ being the totally antisymmetric symbol on $\mathbb{R}^{1,2}$, normalized to $\epsilon_{012} = +1$. Note that this is the same ansatz for $\hat{\phi}$ and \hat{H} as in [24] and that the system of [25] can be recovered by setting \hat{H}_{yuv} and ℓ to zero.

We now proceed by introducing a G_2 structure on the seven-dimensional manifold $X_7 = \mathbb{R} \times X_6$ with metric

$$g_7 = e^{2\Delta(x^u)} dx^3 dx^3 + g_{uv}(x^m) dx^u dx^v. \quad (2.6)$$

The G_2 structure form $\varphi \in \Lambda^3(X_7)$ and its seven-dimensional Hodge dual $\Phi := *_7 \varphi \in \Lambda^4(X_7)$ are constructed using the seven-dimensional gamma matrices Γ_m and the spinor η , which is parallel with respect to ${}^\nabla$,

$$\varphi_{mnp} = -i\eta^\dagger \Gamma_{mnp} \eta, \quad \Phi_{mnpq} = \eta^\dagger \Gamma_{mnpq} \eta. \quad (2.7)$$

The gamma matrices Γ_m are taken to satisfy $\{\Gamma_m, \Gamma_n\} = 2(g_7)_{mn}$, and we define a totally anti-symmetrized product of gamma matrices as $\Gamma_{m_1 \dots m_p} := \Gamma_{[m_1} \cdots \Gamma_{m_p]}$. The first two

equations in (2.1) then imply the following relations [24, 25],

$$\begin{aligned}
 d_7 \varphi &= 2d_7 \hat{\phi} \wedge \varphi - *_7 \hat{H} - \ell \Phi , \\
 d_7 \Phi &= 2d_7 \hat{\phi} \wedge \Phi , \\
 *_7 d_7 \hat{\phi} &= -\frac{1}{2} \hat{H} \wedge \varphi , \\
 0 &= \frac{1}{2} *_7 \ell - \hat{H} \wedge \Phi .
 \end{aligned} \tag{2.8}$$

Here, d_7 and $*_7$ are the differential and the Hodge star defined on X_7 .

Taking into account the decomposition $X_7 = \mathbb{R} \times X_6$, we can rewrite these equations in terms of an $SU(3)$ structure defined on X_6 and the domain wall direction. First, we decompose η into two six-dimensional spinors of definite chirality,

$$\eta = \frac{1}{\sqrt{2}}(\eta_+ + \eta_-) . \tag{2.9}$$

An $SU(3)$ structure on X_6 is uniquely specified via a real 2-form J and a complex 3-form $\Omega = \Omega_+ + i\Omega_-$, which are defined for every fixed value of y using the spinors η_{\pm} [24, 25],

$$\Omega_{uvw} = \eta_+^\dagger \gamma_{uvw} \eta_- , \quad J_{uv} = \mp \eta_\pm^\dagger \gamma_{uv} \eta_\pm , \tag{2.10}$$

where the γ_u are gamma matrices on X_6 satisfying $\{\gamma_u, \gamma_v\} = 2g_{uv}$. The forms (J, Ω) obey the following relations,

$$J \wedge \Omega = 0 , \quad \frac{1}{3!} J \wedge J \wedge J = \frac{i}{8} \Omega \wedge \bar{\Omega} = *1 , \quad *_7 J = \frac{1}{2} J \wedge J , \quad *\Omega_\pm = \pm \Omega_\mp , \tag{2.11}$$

where $*$ is the six-dimensional Hodge star with respect to the metric $g_{uv}(x^m)$. The Hodge star $*$ is in our conventions related to $*_7$ via

$$*_7 \omega_p^{(6)} = e^\Delta (*\omega_p^{(6)}) \wedge dy , \quad *_7(dy \wedge \omega_p^{(6)}) = e^{-\Delta} *\omega_p^{(6)} . \tag{2.12}$$

Here, $\omega_p^{(6)}$ is a p -form with legs only in the X_6 directions.

The relation between the G_2 structure (φ, Φ) and the $SU(3)$ structure (J, Ω) can be expressed as [24–26]

$$\begin{aligned}
 \varphi &= e^\Delta dy \wedge J + \Omega_- , \\
 \Phi &= e^\Delta dy \wedge \Omega_+ + \frac{1}{2} J \wedge J ,
 \end{aligned} \tag{2.13}$$

where the prefactor e^Δ is a consequence of the metric ansatz (2.2).

Substituting the decomposition (2.13) into (2.8), one obtains

$$\begin{aligned}
 dJ &= e^{-\Delta} \Omega'_- - 2e^{-\Delta} \hat{\phi}' \Omega_- + 2d\hat{\phi} \wedge J - J \wedge d\Delta - *H + \ell \Omega_+ , \\
 J \wedge dJ &= J \wedge J \wedge d\hat{\phi} , \\
 d\Omega_+ &= e^{-\Delta} J \wedge J' - e^{-\Delta} \hat{\phi}' J \wedge J + 2d\hat{\phi} \wedge \Omega_+ + \Omega_+ \wedge d\Delta ,
 \end{aligned}$$

$$\begin{aligned}
 d\Omega_- &= 2d\hat{\phi} \wedge \Omega_- - e^{-\Delta} * H_y - \frac{1}{2}\ell J \wedge J, \\
 *d\hat{\phi} &= \frac{1}{2}e^{-\Delta} H_y \wedge \Omega_- - \frac{1}{2}H \wedge J, \\
 *\hat{\phi}' &= -\frac{1}{2}e^{\Delta} H \wedge \Omega_-, \\
 0 &= \frac{1}{2} * \ell - \Omega_+ \wedge H - \frac{1}{2}e^{-\Delta} H_y \wedge J \wedge J.
 \end{aligned} \tag{2.14}$$

Here, a prime denotes the derivative with respect to the coordinate y , and d is the exterior derivative on the six-dimensional manifold X_6 . The two exterior derivatives d_7 and d are related via

$$d_7\omega = d\omega + dy \wedge \omega' \tag{2.15}$$

for some p -form ω . The ten-dimensional NS 3-form \hat{H} is taken to decompose into the following parts, respecting (2.5),

$$\hat{H} = H + dy \wedge H_y + \ell \text{vol}_{\mathbb{R}^{1,2}}, \tag{2.16}$$

with

$$H = \frac{1}{3!} H_{uvw} dx^u \wedge dx^v \wedge dx^w \quad \text{and} \quad H_y = \frac{1}{2!} H_{yuv} dx^u \wedge dx^v \tag{2.17}$$

having legs solely in the X_6 directions. We also define the volume form of $\mathbb{R}^{1,2}$ as $\text{vol}_{\mathbb{R}^{1,2}} := \frac{1}{3!} \epsilon_{\alpha\beta\gamma} dx^\alpha \wedge dx^\beta \wedge dx^\gamma$.

Note that (2.14) may be regarded as a generalization of the Hitchin flow equations [27]. This is a common situation in four-dimensional BPS domain-wall solutions of ten-dimensional supergravity theories [24, 25, 28–30]. In the absence of \hat{H} , $\hat{\phi}$ and Δ , the system of equations in (2.14) reduces to

$$\begin{aligned}
 J \wedge dJ &= 0, & dJ &= \Omega'_-, \\
 d\Omega_- &= 0, & d\Omega_+ &= J \wedge J',
 \end{aligned} \tag{2.18}$$

which are the original Hitchin flow equations.

The structure forms J and Ω are tightly related to the torsion classes defined as irreducible representations of the torsion $T_{mn}{}^p$ under the stability group $\text{SU}(3)$. A manifold with $\text{SU}(3)$ structure in general has a connection with torsion

$$T_{mn}{}^p \in \Lambda^1 \otimes \mathfrak{so}(6). \tag{2.19}$$

The 1-form index is the upper index of the torsion tensor, while the lower antisymmetric pair of indices $[mn]$ label an element of $\mathfrak{so}(6) = \mathfrak{su}(3) \oplus \mathfrak{su}(3)^\perp$. Decomposing the torsion into irreducible representations of the holonomy group and taking into account that the $\mathfrak{su}(3)$ piece drops out when acting on $\text{SU}(3)$ -invariant forms, we obtain the intrinsic torsion [2]

$$\begin{aligned}
 T_{mn}^0{}^p &\in \Lambda^1 \otimes \mathfrak{su}(3)^\perp = (\mathbf{3} \oplus \bar{\mathbf{3}}) \otimes (\mathbf{1} \oplus \mathbf{3} \oplus \bar{\mathbf{3}}) \\
 &= (\mathbf{1} \oplus \mathbf{1}) \oplus (\mathbf{8} \oplus \mathbf{8}) \oplus (\bar{\mathbf{6}} \oplus \mathbf{6}) \oplus 2(\mathbf{3} \oplus \bar{\mathbf{3}}).
 \end{aligned} \tag{2.20}$$

W_1
 W_2
 W_3
 W_4, W_5

The tensors W_1, \dots, W_5 are the five torsion classes that appear in the derivatives of the structure forms,

$$\begin{aligned} dJ &= -\frac{3}{2} \text{Im}(W_1 \bar{\Omega}) + W_4 \wedge J + W_3 , \\ d\Omega &= W_1 J \wedge J + W_2 \wedge J + \bar{W}_5 \wedge \Omega . \end{aligned} \quad (2.21)$$

We now depart from the general discussion and focus on nearly Kähler manifolds, which support instanton connections of the type found in [1] and are defined by the following condition on the torsion classes,

$$W_2 = W_3 = W_4 = W_5 = 0 , \quad \text{while} \quad W_1 = W_1^+ + iW_1^- \quad (2.22)$$

is the only non-zero contribution to the intrinsic torsion. In addition, we set

$$A = 0 \quad \text{and} \quad \Delta = 0 . \quad (2.23)$$

The system of equations in (2.14) is then solved by an NS 3-form flux and a dilaton of the form [24],

$$\begin{aligned} \hat{H} &= -\frac{1}{2} \phi' \Omega_+ + \left(\frac{3}{2} W_1^- + \frac{7}{8} \ell \right) \Omega_- - (2W_1^- + \ell) J \wedge dy + \ell \text{vol}_{\mathbb{R}^{1,2}} , \\ \hat{\phi} &= \phi(y) , \end{aligned} \quad (2.24)$$

provided the structure forms J and Ω satisfy the following flow and structure equations,

$$\begin{aligned} J' &= (W_1^+ + \phi') J , & dJ &= -\frac{3}{2} W_1^- \Omega_+ + \frac{3}{2} W_1^+ \Omega_- , \\ \Omega'_- &= -\left(3W_1^- + \frac{15}{8} \ell \right) \Omega_+ + \frac{3}{2} (W_1^+ + \phi') \Omega_- , & d\Omega &= W_1 J \wedge J . \end{aligned} \quad (2.25)$$

By acting with a y -derivative on the second equation in (2.11), one also learns that

$$\Omega'_+ = \frac{3}{2} (W_1^+ + \phi') \Omega_+ + \alpha(y) \Omega_- , \quad (2.26)$$

with some as yet undetermined function $\alpha(y)$. The expressions (2.24)–(2.26) represent the most general solution of the first two BPS equations in (2.1) with the general metric ansatz (2.2) and $A = \Delta = 0$ on a nearly Kähler manifold X_6 .

Before constructing order α' solutions, we shall first discuss the zeroth order case. We have already solved the first two BPS equations in (2.1). In addition, the third BPS equation in (2.1) is solved by $\hat{F} = 0$. In order to have a full heterotic supergravity solution, we also need to check that the Bianchi identity and the time-like components of the equations of motion are satisfied. The latter leads to the condition $\ell = 0$, as will be shown in more detail in section 4.3. The Bianchi identity at the zeroth order in α' simply becomes $\hat{d}\hat{H} = 0$. From this condition, we obtain the following set of equations,

$$0 = \phi' W_1^+ - 3(W_1^-)^2 , \quad (2.27)$$

$$0 = \phi'' + \frac{3}{2} (\phi')^2 + \frac{13}{2} \phi' W_1^+ , \quad (2.28)$$

$$0 = \phi' \alpha - 3(W_1^-)' - \frac{21}{2} W_1^+ W_1^- - \frac{9}{2} \phi' W_1^- . \quad (2.29)$$

We can immediately read off two special solutions

$$1. \quad \phi = \text{const.}, \quad W_1^+ = \text{any}, \quad W_1^- = 0, \quad \alpha = \text{any}, \quad (2.30)$$

$$2. \quad \phi = \frac{2}{3} \log(ay + b), \quad W_1^+ = 0, \quad W_1^- = 0, \quad \alpha = 0, \quad (2.31)$$

where a, b are integration constants and ‘any’ means a free function. The first case corresponds to a nearly Kähler geometry with constant dilaton and vanishing NS 3-form flux. The second case is Calabi-Yau with flux. Both solutions are contained in [24] as special cases. This concludes our analysis of the zeroth order case, and we shall turn to the construction of order α' solutions.

3 Yang-Mills sector

3.1 Yang-Mills instantons on $\mathbb{R} \times X_6$

In this section, we review the construction of Yang-Mills instantons à la Harland and Nölle [1]. In their terminology, an instanton is a solution of $\hat{F} \cdot \epsilon = 0$, which is the third BPS equation in (2.1). At the zeroth order in α' , one may simply set $\hat{F} = 0$ and ignore the Yang-Mills sector altogether. This is consistent, since the coupling of \hat{F} to the other supergravity fields only starts to arise at linear order in α' . Since our goal is to construct order α' solutions, we need a non-trivial \hat{F} .

We will study the instanton equation on the manifold $X_7 = \mathbb{R} \times X_6$ with ‘ h -cone’ metric¹

$$g_7 = dy^2 + (h(y))^2 \tilde{g}, \quad (3.1)$$

where \tilde{g} is a fixed (that is y -independent) nearly Kähler metric on X_6 with components given by $\tilde{g}_{uv}(x^u)$, and $h(y)$ is a warp factor. Note that this metric is further restricted as compared to the g_7 introduced in the previous section. We henceforth take g_7 to have the form (3.1).

The orthonormal frame on X_7 is given by $\{\sigma^{\underline{m}}\} = \{dy, h e^{\underline{u}}\}$ with $\underline{m} = \underline{3}, \underline{4}, \dots, \underline{9}$ and $\underline{u} = \underline{4}, \underline{5}, \dots, \underline{9}$. Here, $\{e^{\underline{u}}\} = \{e^{\underline{u}}_a dx^a\}$ is an orthonormal frame on X_6 satisfying $e^{\underline{u}}_a e^{\underline{v}}_b \delta_{\underline{u}\underline{v}} = g_{ab}$. Associated to the y -independent metric \tilde{g} , there is a static (that is y -independent) SU(3) structure on X_6 defined in terms of a real 2-form \tilde{J} and a complex 3-form $\tilde{\Omega}$. The orthonormal frame $\{e^{\underline{u}}\}$ on X_6 can be arranged such that \tilde{J} and $\tilde{\Omega}$ take the following standard form,

$$\begin{aligned} \tilde{J} &= e^{\underline{4}} \wedge e^{\underline{5}} + e^{\underline{6}} \wedge e^{\underline{7}} + e^{\underline{8}} \wedge e^{\underline{9}}, \\ \tilde{\Omega} &= (e^{\underline{4}} + i e^{\underline{5}}) \wedge (e^{\underline{6}} + i e^{\underline{7}}) \wedge (e^{\underline{8}} + i e^{\underline{9}}). \end{aligned} \quad (3.2)$$

These forms satisfy $d\tilde{\Omega}_+ = 2\tilde{J} \wedge \tilde{J}$ and $d\tilde{J} = 3\tilde{\Omega}_-$, showing that the manifold X_6 is indeed nearly Kähler with $\tilde{W}_1^+ = 2$ and $\tilde{W}_1^- = 0$. In addition, they obey the equations in (2.11) with tildes everywhere.

¹The metric (3.1) may be regarded as a generalized cone metric. It reduces to the standard cone metric on $\mathbb{R}_+ \times X_6$ upon setting $h(y) = y$.

The instanton equation $\hat{F} \cdot \epsilon = 0$ on X_7 reduces to

$$*_7 \hat{F} = -(*_7 Q) \wedge \hat{F} \quad \text{where} \quad Q = h^3 dy \wedge \tilde{\Omega}_+ + \frac{1}{2} h^4 \tilde{J} \wedge \tilde{J}. \quad (3.3)$$

Via the coordinate redefinition, assuming $h \geq 0$ (see footnote 3),

$$dy = e^{f(\tau)} d\tau \quad \text{where} \quad e^{f(\tau)} = h(y(\tau)), \quad (3.4)$$

the metric (3.1) transforms into

$$g_7 = e^{2f} g_Z \quad \text{with} \quad g_Z = d\tau^2 + \tilde{g}, \quad (3.5)$$

where g_Z is the metric on the cylinder. Since (3.3) is conformally invariant and the metric (3.1) is conformal to the cylinder metric, instantons on the cylinder will also solve (3.3). The instanton equation on the cylinder is

$$*_Z \hat{F} = -(*_Z Q_Z) \wedge \hat{F} \quad \text{where} \quad Q_Z = d\tau \wedge \tilde{\Omega}_+ + \frac{1}{2} \tilde{J} \wedge \tilde{J}, \quad (3.6)$$

and $*_Z$ is the Hodge star with respect to the cylinder metric g_Z .

To solve (3.6), we use the same ansatz for the gauge connection as in [1], namely

$${}^A \hat{\nabla} = {}^{\text{can}} \nabla + \psi(\tau) e^u I_{\underline{u}}, \quad (3.7)$$

where ${}^{\text{can}} \nabla$ is the canonical connection on X_6 defined by

$${}^{\text{can}} \omega_u^{\underline{v}}{}_{\underline{w}} = {}^{\text{LC}} \omega_u^{\underline{v}}{}_{\underline{w}} + \frac{1}{2} (\tilde{\Omega}_+)_{\underline{w}\underline{v}\underline{u}} e^u, \quad (3.8)$$

and the matrices $I_{\underline{u}}$, generating the orthogonal complement of $\mathfrak{su}(3)$ in $\mathfrak{g}_2 \subset \mathfrak{so}(7)$ are defined by the following relations,

$$(I_{\underline{u}})^{\underline{3}}{}_{\underline{v}} = -(I_{\underline{u}})^{\underline{v}}{}_{\underline{3}} = -\delta_{\underline{v}}^{\underline{u}}, \quad (I_{\underline{u}})^{\underline{w}}{}_{\underline{v}} = -\frac{1}{2} (\tilde{\Omega}_+)_{\underline{u}\underline{v}\underline{w}}. \quad (3.9)$$

Together with the generators of $\mathfrak{su}(3)$, denoted $(I_i)^{\underline{u}}{}_{\underline{v}}$, they form a basis of the Lie algebra \mathfrak{g}_2 . The curvature 2-form $\hat{F} = \frac{1}{2} [{}^A \hat{\nabla}, {}^A \hat{\nabla}]$ of the gauge connection (3.7) becomes [1]

$$\hat{F} = {}^{\text{can}} R + \frac{1}{2} \psi^2 f_{\underline{u}\underline{v}}^i e^u \wedge e^v I_i + \dot{\psi} d\tau \wedge e^u I_{\underline{u}} + \frac{1}{2} (\psi - \psi^2) (\tilde{\Omega}_+)_{\underline{u}\underline{v}\underline{w}} e^v \wedge e^w I_{\underline{u}} =: \mathcal{F}(\psi), \quad (3.10)$$

where $f_{\underline{u}\underline{v}}^i$ is a structure constant appearing in the Lie algebra commutator

$$[I_{\underline{u}}, I_{\underline{v}}] = f_{\underline{u}\underline{v}}^i I_i + f_{\underline{u}\underline{v}}^w I_{\underline{w}}, \quad (3.11)$$

and a dot denotes a derivative with respect to τ . It was shown in [1] that such an \hat{F} solves the instanton equation (3.6) if and only if the function $\psi(y)$ satisfies the ‘kink equation’

$$\dot{\psi} = 2\psi(\psi - 1). \quad (3.12)$$

This equation has two fixed points, $\psi = 0$ and $\psi = 1$, which correspond to the canonical connection ${}^{\text{can}} \nabla$ and the Levi-Civita connection ${}^{\text{LC}} \nabla$ on X_7 , respectively. There is also

a non-constant solution. It interpolates between the two fixed points and is given by the kink function

$$\psi(\tau) = \frac{1}{2} (1 - \tanh[\tau - \tau_0]) . \quad (3.13)$$

The integration constant τ_0 fixes the position of the instanton in the τ direction. In terms of the original variables y and $h(y)$, the kink equation becomes

$$h(y) \psi'(y) = 2 \psi(y) (\psi(y) - 1) . \quad (3.14)$$

It has the same fixed points, $\psi = 0$ and $\psi = 1$, as (3.12). The non-constant solution is formally given by

$$\psi(y) = \frac{1}{2} (1 - \tanh[\tau(y) - \tau_0]) , \quad (3.15)$$

with $\tau(y)$ determined by (3.4).

Heterotic supergravity contains another curvature 2-form besides \hat{F} , namely \tilde{R} . As explained in section 1, we adopt the supergravity point of view for the purpose of this paper, which implies the instanton equation also for \tilde{R} , that is

$$\tilde{R} \cdot \epsilon = 0 . \quad (3.16)$$

Given that we have found an explicit instanton solution for the ansatz (3.7), we will make the same ansatz also for $\tilde{\nabla}$. Each connection, however, comes equipped with its own independent scalar function ψ . To distinguish between the two, we put

$$\tilde{R} = \mathcal{F}(\psi_1) \quad \text{and} \quad \hat{F} = \mathcal{F}(\psi_2) , \quad (3.17)$$

where $\mathcal{F}(\psi)$ was defined in (3.10). The precise choices for $\psi_{1,2}$ will be made later.

3.2 Bianchi identity

The Green-Schwarz anomaly cancellation condition for the heterotic string at order α' can be written as a Bianchi identity for the NS 3-form \hat{H} , namely [9, 13]

$$\hat{d}\hat{H} = \frac{\alpha'}{4} \text{Tr}(\hat{F} \wedge \hat{F} - \tilde{R} \wedge \tilde{R}) . \quad (3.18)$$

With the results from the previous subsection, one may explicitly evaluate the right-hand side of this equation. First, one has

$$\hat{d}\hat{H} = -\frac{\alpha'}{4} \text{Tr}(\mathcal{F}(\psi_1) \wedge \mathcal{F}(\psi_1) - \mathcal{F}(\psi_2) \wedge \mathcal{F}(\psi_2)) . \quad (3.19)$$

Inserting (3.10) and using the kink equation (3.14), we can express the Bianchi identity in the following form,

$$\hat{d}\hat{H} = -\frac{\alpha'}{4} \left[3h((\psi'_1)^2 - (\psi'_2)^2) dy \wedge \tilde{\Omega}_+ + 2(\psi_1^2(2\psi_1 - 3) - \psi_2^2(2\psi_2 - 3)) \tilde{J} \wedge \tilde{J} \right] , \quad (3.20)$$

which will turn out to be useful in the following sections.

4 Warped nearly Kähler domain wall

4.1 Static versus dynamic SU(3) structures on X_6

In section 2, we began our general discussion by assuming a dynamic, that is y -dependent, SU(3) structure on (X_6, g) characterized by a pair of forms (J, Ω) . This was subsequently specialized in section 3 to the case where $g = (h(y))^2 \tilde{g}$ with a static, that is y -independent, SU(3) structure on (X_6, \tilde{g}) with the forms $(\tilde{J}, \tilde{\Omega})$.

Given that a pair of SU(3) structure forms uniquely specifies a metric and the relation

$$g = (h(y))^2 \tilde{g} , \quad (4.1)$$

it is clear that the two pairs (J, Ω) and $(\tilde{J}, \tilde{\Omega})$ are not unrelated. Indeed, the Hodge star satisfies $*\omega_p = h^{6-2p} \tilde{*}\omega_p$ for a p -form ω_p on X_6 , and, together with $\frac{1}{3!} J^3 = *1$ as well as $\frac{1}{3!} \tilde{J}^3 = \tilde{*}1$, this implies

$$J = h^2 \tilde{J} . \quad (4.2)$$

The relation between Ω and $\tilde{\Omega}$ is a little more subtle, due to the fact that there can also be a mixing between real and imaginary parts. To parametrize this mixing, we write

$$\Omega_+ = h^3 \cos \beta \tilde{\Omega}_+ + h^3 \sin \beta \tilde{\Omega}_- , \quad \Omega_- = -h^3 \sin \beta \tilde{\Omega}_+ + h^3 \cos \beta \tilde{\Omega}_- , \quad (4.3)$$

with a y -dependent mixing angle $\beta \in [0, 2\pi)$. A shift of $\beta \rightarrow \beta + \pi$ can be compensated by a sign flip of h , and so we may restrict ourselves to $\beta \in [0, \pi)$. The chosen parametrization automatically guarantees that

$$\begin{aligned} \frac{i}{8} \tilde{\Omega} \wedge \tilde{\Omega} &= \tilde{*}1 & \implies & \frac{i}{8} \Omega \wedge \bar{\Omega} = *1 , & \text{and} \\ \tilde{*}\tilde{\Omega}_\pm &= \pm \tilde{\Omega}_\mp & \implies & *\Omega_\pm = \pm \Omega_\mp , \end{aligned} \quad (4.4)$$

which is required in order to be compatible with our SU(3) structure conventions as formulated in (2.11). Finally, comparing the expressions for dJ and $d\Omega$ with the tilded versions in section 3.1, we learn that

$$W_1^+ = 2h^{-1} \cos \beta \quad \text{and} \quad W_1^- = -2h^{-1} \sin \beta . \quad (4.5)$$

Thus, mixing in (4.3) occurs unless either W_1^+ or W_1^- vanish. Setting both W_1^+ and W_1^- to zero and thereby reducing to a Calabi-Yau geometry is not possible.²

Upon inserting the above expressions into the flow equations in (2.25), it follows that

$$h' = \cos \beta + \frac{1}{2} h \phi' , \quad (4.6)$$

$$0 = h^2 \left(h \beta' + 6 \sin \beta - \frac{15}{8} \ell h \right) . \quad (4.7)$$

Moreover, the as yet unknown coefficient function $\alpha(y)$ in (2.26) is now fixed as

$$\alpha(y) = 3W_1^- + \frac{15}{8} \ell = -6h^{-1} \sin \beta + \frac{15}{8} \ell . \quad (4.8)$$

²We are interested in solutions where (X_6, g) is compact. Here and in the following, we thus consider only finite h and exclude the possibility of taking the decompactification limit $h \rightarrow \pm\infty$.

The expression (2.24) for the NS 3-form \hat{H} in terms of $(\tilde{J}, \tilde{\Omega})$ becomes

$$\begin{aligned}\hat{H} = & h^2 \left[-\frac{1}{2} h \phi' \cos \beta + 3 \sin^2 \beta - \frac{7}{8} \ell h \sin \beta \right] \tilde{\Omega}_+ \\ & + h^2 \left[-\frac{1}{2} h \phi' \sin \beta - 3 \sin \beta \cos \beta + \frac{7}{8} \ell h \cos \beta \right] \tilde{\Omega}_- \\ & + h (4 \sin \beta - \ell h) \tilde{J} \wedge dy + \ell \text{vol}_{\mathbb{R}^{1,2}} .\end{aligned}\quad (4.9)$$

It is instructive to pause here and reflect on what we have achieved so far. Provided the coupled ordinary differential equations (4.6)–(4.7) involving the scalar functions h , β , ℓ and ϕ are satisfied, we have a solution of the first two BPS equations in (2.1) for the metric ansatz

$$\hat{g} = \eta_{\alpha\beta} dx^\alpha dx^\beta + dx^3 dx^3 + (h(x^3))^2 \tilde{g}_{uv}(x^w) dx^u dx^v , \quad (4.10)$$

and the restrictions in (2.5). The third BPS equation in (2.1) is solved by the instanton construction presented in section 3.1. However, we still need to ensure that the Bianchi identity is satisfied and check that the time-like components of the equations of motion are obeyed. These issues will be addressed in the next two subsections.

4.2 Embedding of the instanton in the ten-dimensional solution

In order to embed the instanton solution of section 3.1 into a fully ten-dimensional solution, we need to impose the Bianchi identity (3.18). In section 3.2, we have already computed the right-hand side of the Bianchi identity. The left-hand side can be further specified by applying a ten-dimensional exterior derivative \hat{d} on the expression (4.9) for \hat{H} ,

$$\begin{aligned}\hat{d}\hat{H} = & \left\{ \left[-\frac{1}{2} h^3 \phi' \sin \beta - 3 h^2 \sin \beta \cos \beta + \frac{7}{8} \ell h^3 \cos \beta \right]' - 3 h (4 \sin \beta - \ell h) \right\} dy \wedge \tilde{\Omega}_- \\ & + \left[-\frac{1}{2} h^3 \phi' \cos \beta + 3 h^2 \sin^2 \beta - \frac{7}{8} \ell h^3 \sin \beta \right]' dy \wedge \tilde{\Omega}_+ \\ & + 2 \left[-\frac{1}{2} h^3 \phi' \cos \beta + 3 h^2 \sin^2 \beta - \frac{7}{8} \ell h^3 \sin \beta \right] \tilde{J} \wedge \tilde{J} \\ & + \ell' dy \wedge \text{vol}_{\mathbb{R}^{1,2}} .\end{aligned}\quad (4.11)$$

Comparing with (3.20), we obtain the following additional conditions on the scalar functions h , β , ℓ and ϕ , now also coupled to the instanton solutions $\psi_{1,2}$ as given in (3.13),

$$h^3 \phi' \cos \beta - 6 h^2 \sin^2 \beta + \frac{7}{4} \ell h^3 \sin \beta = \frac{\alpha'}{2} (\psi_1^2 (2\psi_1 - 3) - \psi_2^2 (2\psi_2 - 3)) , \quad (4.12)$$

$$\left[h^3 \phi' \sin \beta + 6 h^2 \sin \beta \cos \beta - \frac{7}{4} \ell h^3 \cos \beta \right]' = -6 h (4 \sin \beta - \ell h) , \quad (4.13)$$

$$\ell' = 0 . \quad (4.14)$$

From the $dy \wedge \tilde{\Omega}_+$ term in (4.11), one obtains also the y -derivative of (4.12). This, however, yields no further condition and is thus omitted. The relation (4.13) may be re-written in

the form

$$\sin \beta \left(h^2 \phi'' + \frac{3}{2} (h\phi')^2 + 27h\phi' \cos \beta + \frac{105}{32} \ell^2 h^2 - \frac{111}{4} \ell h \sin \beta + 12 \sin^2 \beta + 48 \right) - \frac{3}{4} \ell h^2 \phi' \cos \beta = 0, \quad (4.15)$$

after using (4.6), (4.7) and (4.14).

4.3 Equations of motion

The equations of motion of heterotic supergravity up to and including terms of order α' read

$$\begin{aligned} \hat{R}_{\mu\nu} + 2(\hat{\nabla} \hat{d}\hat{\phi})_{\mu\nu} - \frac{1}{4} \hat{H}_{\kappa\lambda\mu} \hat{H}_{\nu}{}^{\kappa\lambda} + \frac{\alpha'}{4} \left[\tilde{R}_{\mu\kappa\lambda\sigma} \tilde{R}_{\nu}{}^{\kappa\lambda\sigma} - \text{tr} \left(\hat{F}_{\mu\kappa} \hat{F}_{\nu}{}^{\kappa} \right) \right] &= 0, \\ \hat{R} + 4\hat{\Delta}\hat{\phi} - 4|\hat{d}\hat{\phi}|^2 - \frac{1}{2} |\hat{H}|^2 + \frac{\alpha'}{4} \text{tr} \left[|\tilde{R}|^2 - |\hat{F}|^2 \right] &= 0, \\ e^{2\hat{\phi}} \hat{d} \hat{*} (e^{-2\hat{\phi}} \hat{F}) + \hat{A} \wedge \hat{*} \hat{F} - \hat{*} \hat{F} \wedge \hat{A} + \hat{*} \hat{H} \wedge \hat{F} &= 0, \\ \hat{d} \hat{*} e^{-2\hat{\phi}} \hat{H} &= 0. \end{aligned} \quad (4.16)$$

Here, $\hat{R}_{\mu\nu}$ and \hat{R} are the Ricci tensor and the scalar curvature, respectively. They are computed from the full ten-dimensional metric (4.10). The gauge field \hat{A} corresponds to the curvature $\hat{F} = \mathcal{F}(\psi_2)$, and its components can be read off from

$${}^A \hat{\nabla} = {}^{\text{can}} \nabla + \psi_2(y) e^{\underline{u}} I_{\underline{u}} = \hat{d} + \hat{A}. \quad (4.17)$$

We will, however, not need the explicit form of this field. Finally, we define $|\omega|^2 := \frac{1}{p!} \omega_{\mu_1 \dots \mu_p} \omega^{\mu_1 \dots \mu_p}$ for a p -form ω , and we note that the Einstein equation has been simplified by means of the dilaton equation.

Since we adopted the supergravity point of view on the curvature \tilde{R} , we do not need to verify explicitly all the equations of motion. The precise implication for the equations of motion following from the BPS equations and Bianchi identity is a somewhat subtle point [13, 24, 31, 32] (for recent discussions, see also [33–36]). We remark that ansätze for the connections and various other assumptions, such as the considered order in the α' expansion, differ in the literature, and this effects the conclusions that are drawn. For the purpose of this paper, we shall follow a conservative strategy and assume that, for our set-up, the BPS equations together with the Bianchi identity and the *time-like* components of the equations of motion imply the remaining components of the equations of motion.

The time-like components of the Yang-Mills equation are trivially satisfied by the ansatz, since the time-like components $\hat{F}_{0\mu}$ of the corresponding field strength are identically zero. The same is true for the mixed (0μ) -components of the Einstein equations where $\mu \neq 0$. Hence, we are left with the following two equations,

$$\hat{R}_{00} + 2(\hat{\nabla} \hat{d}\hat{\phi})_{00} - \frac{1}{4} \hat{H}_{0\mu\nu} \hat{H}_0{}^{\mu\nu} = 0, \quad (4.18)$$

$$\hat{\nabla}_{\mu} \left(e^{-2\hat{\phi}} \hat{H}^{\mu\nu 0} \right) = 0. \quad (4.19)$$

For our field ansatz, (4.19) is satisfied trivially, whereas (4.18) implies

$$\ell = 0 . \quad (4.20)$$

This condition eliminates any flux that has legs in the domain-wall world-volume directions.

4.4 Summary of the system of equations

For the warped nearly Kähler domain wall considered in this paper, we have obtained in the previous sections a system of coupled non-linear ordinary differential equations for the scalar functions h , β and ϕ . This system of equations is a consequence of consistently embedding the instanton on $\mathbb{R} \times X_6$ into a full-fledged heterotic supergravity solution with NS 3-form flux given by (4.9). Before we study explicit solutions, it is beneficial to first collect and summarize the complete system of equations. This shall then serve as the central point of reference for the remainder of this paper.

Having set $\ell = 0$ in accordance with the result of the previous subsection, the full set of equations then reads as follows,

$$h' = \cos \beta + \frac{1}{2} h \phi' , \quad (4.21)$$

$$0 = h^2 (h \beta' + 6 \sin \beta) , \quad (4.22)$$

$$0 = h^3 \phi' \cos \beta - 6 h^2 \sin^2 \beta - \frac{\alpha'}{2} (\psi_1^2 (2\psi_1 - 3) - \psi_2^2 (2\psi_2 - 3)) , \quad (4.23)$$

$$0 = \sin \beta \left(h^2 \phi'' + \frac{3}{2} (h \phi')^2 + 27 h \phi' \cos \beta + 12 \sin^2 \beta + 48 \right) , \quad (4.24)$$

$$h \psi_1' = 2 \psi_1 (\psi_1 - 1) , \quad (4.25)$$

$$h \psi_2' = 2 \psi_2 (\psi_2 - 1) . \quad (4.26)$$

This is an *a priori* over-determined system since there are six equations for the five scalar functions h , β , ϕ , ψ_1 and ψ_2 . It is supplemented by the determination of the torsion class W_1 of the warped six-dimensional space (X_6, g) via $W_1 = 2h^{-1} e^{-i\beta}$.

A solution obeying (4.21)–(4.26) guarantees that the corresponding ten-dimensional fields \hat{g} , \hat{H} , $\hat{\phi}$ and ${}^A\hat{\nabla}$ satisfy the BPS equations (2.1), the Bianchi identity (3.18) and, in turn, also the equations of motion (4.16). Since the connection ${}^{-}\hat{\nabla}$ determined by the NS 3-form flux (4.9) preserves one parallel spinor ϵ as in (2.4), we are dealing with an $\mathcal{N}=1/2$ solution from the point of view of $\mathbb{R}^{1,3}$ spanned by the non-compact coordinates $\{x^0, x^1, x^2, x^3\}$.

An analytic solution to the highly complicated system of equations (4.21)–(4.26) would be too much to ask for. Instead, one may concentrate on special cases and hope to find analytic or numerical solutions there. An obvious specialization is to turn off either W_1^+ or W_1^- .

The first case, that is $W_1^+ = 0$, can be dealt with very quickly. Setting $W_1^+ = 0$ and assuming $h < \infty$ implies $\cos \beta = 0$. However, (4.22) then immediately leads to $h = 0$, which causes the metric to be ill-defined. We thus conclude that a well-defined solution requires $W_1^+ \neq 0$.

Fortunately, the second case, that is $W_1^- = 0$, turns out to be more fruitful. It will be the subject of the next section.

5 Explicit solutions with $W_1^- = 0$

In this section, we investigate the solution of (4.21)–(4.26) for the special case that $W_1^- = 0$. For $h < \infty$ this implies $\sin \beta = 0$, hence $\cos \beta = 1$. (Here, we are restricting to $\beta \in [0, \pi)$, without loss of generality, as explained below (4.3).) Therefore, the relation (4.3) involves no mixing,

$$\Omega_+ = h^3 \tilde{\Omega}_+ \quad \text{and} \quad \Omega_- = h^3 \tilde{\Omega}_- . \quad (5.1)$$

The system of equations in (4.21)–(4.26) then reduces to

$$h' = 1 + \frac{1}{2} h \phi' , \quad (5.2)$$

$$h^3 \phi' = \frac{\alpha'}{2} (\psi_1^2 (2\psi_1 - 3) - \psi_2^2 (2\psi_2 - 3)) , \quad (5.3)$$

$$h \psi_1' = 2 \psi_1 (\psi_1 - 1) , \quad (5.4)$$

$$h \psi_2' = 2 \psi_2 (\psi_2 - 1) , \quad (5.5)$$

and the NS 3-form flux is simply given by

$$\hat{H} = -\frac{1}{2} h^3 \phi' \tilde{\Omega}_+ = -\frac{\alpha'}{4} (\psi_1^2 (2\psi_1 - 3) - \psi_2^2 (2\psi_2 - 3)) \tilde{\Omega}_+ . \quad (5.6)$$

We remark that there are no terms containing $\tilde{\Omega}_-$ and $\tilde{J} \wedge dy$. Indeed, in the absence of ℓ , those contributions are precisely measured by W_1^- , which is taken to vanish here. Hence, the NS 3-form flux is completely internal, that is $H_y = 0$. The solution then belongs to the same class of half-flat constructions as those obtained in [23], given the condition on the dilaton $d\phi = 0$. The precise forms of the solutions, however, depend on the instanton configurations in question. The relation to solutions contained in the existing literature will be clarified below.

Following [1, 22], we may integrate (5.2)–(5.5) after performing a combined coordinate and function redefinition $y \rightarrow \tau$, $h(y) \rightarrow f(\tau)$ of the form³

$$dy = e^{f(\tau)} d\tau , \quad h(y) = e^{f(\tau(y))} . \quad (5.7)$$

The metric then becomes

$$\hat{g} = \eta_{\alpha\beta} dx^\alpha dx^\beta + e^{2f(\tau)} (d\tau^2 + \tilde{g}_{uv}(x^w) dx^u dx^v) , \quad (5.8)$$

³Note that this transformation requires h to be non-negative. However, for the solutions discussed below this is not always the case. For the cases with $\psi_1 = 0, 1$ and $\psi_2 = 0, 1$, one may solve (5.2)–(5.5) directly and this yields a solution valid for all $y \in \mathbb{R}$. In cases where at least one of the $\psi_{1,2}$ is a non-constant solution of (5.4)–(5.5) and when expressed in terms of y , our solutions are only valid on a half-space ranging from the physical location of the domain wall to infinity. It is then interesting to ask whether and how the solution can be continued across the domain wall. A general argument for the existence of such a continuation for all $y \in \mathbb{R}$ has been given in [22], section 5.6. We will, however, not attempt to answer this question in the present paper.

and (5.2)–(5.5) turn into

$$\dot{\phi} = 2(\dot{f} - 1) , \quad (5.9)$$

$$2(\dot{f} - 1)e^{2(f-\tau)} = \frac{\alpha'}{2}e^{-2\tau} (\psi_1^2(2\psi_1 - 3) - \psi_2^2(2\psi_2 - 3)) , \quad (5.10)$$

$$\dot{\psi}_1 = 2\psi_1(\psi_1 - 1) , \quad (5.11)$$

$$\dot{\psi}_2 = 2\psi_2(\psi_2 - 1) , \quad (5.12)$$

where a dot denotes a derivative with respect to the new coordinate τ . Equation (5.9) implies

$$\phi(\tau) = \phi_0 + 2(f - \tau) , \quad (5.13)$$

with a constant of integration denoted ϕ_0 . Equation (5.10) can be integrated as well, and we find

$$e^{2f} = e^{2(\tau-\tau_0)} + \frac{\alpha'}{4} (\psi_1^2 - \psi_2^2) , \quad (5.14)$$

with some constant of integration denoted τ_0 . The different choices for $\psi_{1,2}$ are either 0, 1 or the interpolating kink solution (3.13) with integration constants $\tau_{1,2}$. This leaves us with a total of 8 different instanton configurations, which are discussed below.

Case 1: $\psi_1 = \psi_2$. All α' corrections cancel out and we revert to the zeroth order solution of [23], section 3.6, that is

$$f = \tau - \tau_0 , \quad \phi = \phi_0 - 2\tau_0 . \quad (5.15)$$

This is shown in figure 1. The NS 3-form flux vanishes, $\hat{H} = 0$, the dilaton is constant, and the ten-dimensional metric

$$\hat{g} = \eta_{\alpha\beta} dx^\alpha dx^\beta + dx^3 dx^3 + (x^3)^2 \tilde{g}_{uv}(x^w) dx^u dx^v \quad (5.16)$$

is the cone metric on X_7 together with the flat Minkowski metric on $\mathbb{R}^{1,2}$ (ignoring the irrelevant integration constant τ_0 , which can always be absorbed by a scale transformation of the coordinate $x^3 := e^{\tau-\tau_0}$).

Case 2: $\psi_1 = 1, \psi_2 = 0$. This case admits a static solution with a linear dilaton,

$$f = f_0 := \frac{1}{2} \log \left(\frac{\alpha'}{4} \right) \quad \text{and} \quad \phi = \phi_0 + 2(f_0 - \tau) . \quad (5.17)$$

The general solution can immediately be stated by inserting the values for $\psi_{1,2}$ into (5.13)–(5.14),

$$e^{2f} = e^{2(\tau-\tau_0)} + \frac{\alpha'}{4} \quad \text{and} \quad e^{\phi-\phi_0} = e^{-2\tau_0} + \frac{\alpha'}{4} e^{-2\tau} . \quad (5.18)$$

This is a special case of [1], section 5.1 (with $\psi = 0$) and of [22], section 5.5 (with $Q_e = 0$, $a = 1$ and taking the decompactification limit in the S^1 direction). An exemplary plot showing the behavior of $f(\tau)$ and $\phi(\tau)$ is shown on the left of figure 2.




Figure 1. Solution for case 1 ($\psi_1 = \psi_2$). This is the trivial solution corresponding to a cone metric with constant dilaton and vanishing NS 3-form flux.

The transformation back to the original variable y is problematic, however. Solving (5.7), with the solution (5.18) inserted, formally yields

$$y(\tau) = \frac{1}{2} \left(\sqrt{4e^{2(\tau-\tau_0)} + \alpha'} - \sqrt{\alpha'} \operatorname{artanh} \left[\sqrt{1 + \frac{4}{\alpha'} e^{2(\tau-\tau_0)}} \right] \right) + y_0. \quad (5.19)$$

For finite τ , the argument of artanh is always greater than one and thus $y(\tau)$ is ill-defined over the reals. One may however directly solve (5.2)–(5.5) for this case. After inserting (5.3) into (5.2) and considering the inverse function $y(h)$, we obtain

$$y - y_0 = h - \frac{\sqrt{\alpha'}}{2} \operatorname{artanh} \left(\frac{2h}{\sqrt{\alpha'}} \right). \quad (5.20)$$

Another inversion, which however cannot be performed explicitly, yields $h(y)$. We see that in terms of the original variables y and $h(y)$, this case is equivalent to the scenario discussed in [23], section 4.5.2.

The graph of $h(y)$ has a kink shape and a zero at the value $y = y_0$, which indicates the location of the domain wall. In addition, (5.20) does not have solutions for all values of y . Instead, y is constrained to lie in the interval $|y - y_0| \leq y_{\max}$. The boundary value y_{\max} can only be determined numerically. It depends on α' and y_0 . As $y \rightarrow \pm y_{\max}$, $h(y)$ limits to $\mp \frac{\sqrt{\alpha'}}{2}$. We note that the scalar curvature for the metric (4.10) given by the following expression,

$$\hat{R} = -12h^{-1}h'' - 30h^{-2}(h')^2 + h^{-2}\tilde{R}, \quad (5.21)$$

diverges at the location of the domain wall, $y = y_0$, and we thus expect the supergravity approximation to break down in the vicinity of the domain wall. In the expression above, \tilde{R} denotes the scalar curvature of the static nearly Kähler metric \tilde{g} on X_6 , which is conventionally normalized to $\tilde{R} = 30$ [37].




Figure 2. Solution for case 2 ($\psi_1 = 1$, $\psi_2 = 0$). The solution is plotted in terms of the new coordinate τ on the left and in terms of the old coordinate y on the right. We have adjusted the free constants to $\alpha' = 64$, $\tau_0 = 0$, $\phi_0 = 3$ and $y_0 = 0$, for concreteness.

The solution for $\phi(y)$ can be implicitly written as

$$\phi(h) = \log \left(\frac{h^2}{4h^2 - \alpha'} \right) + \phi_0, \quad (5.22)$$

where it is understood that the solution for $h(y)$ should be inserted. Taken at face value, this solution is ill-defined as a function of y , even inside the range $|y - y_0| \leq y_{\max}$, since the argument inside the log is negative. This can be cured by using that $\log(-|x|) = \log|x| + i\pi$, and absorbing the second term into a new integration constant $\tilde{\phi}_0 = \phi_0 + i\pi$. The solution for ϕ then reads as follows,

$$\phi(h) = \log \left(\left| \frac{h^2}{4h^2 - \alpha'} \right| \right) + \tilde{\phi}_0. \quad (5.23)$$

As $y \rightarrow \pm y_{\max}$ we have $\phi \rightarrow \infty$, and as $y \rightarrow y_0$ we have $\phi \rightarrow -\infty$. The plots of $h(y)$ and $\phi(y)$, ignoring the integration constant $\tilde{\phi}_0$, are shown on the right of figure 2.

Case 3: $\psi_1 = 0$, $\psi_2 = 1$. Contrary to the previous case, there is no static solution in this case. The general solution (5.13)–(5.14) now becomes

$$e^{2f} = e^{2(\tau - \tau_0)} - \frac{\alpha'}{4}, \quad e^{\phi - \phi_0} = e^{-2\tau_0} - \frac{\alpha'}{4} e^{-2\tau}. \quad (5.24)$$

This case appeared neither in [1] nor in [22]. It should be noted that the expression for $f(\tau)$ following from (5.24) is ill-defined for $\tau < \tau_{\text{dw}}$, where $\tau_{\text{dw}} = \tau_0 + \frac{1}{2} \log(\frac{\alpha'}{4})$ is the location of the domain wall. For $\tau \rightarrow \tau_{\text{dw}}$ from above, the scalar curvature \hat{R} , given in terms of the new variables τ and $f(\tau)$ by⁴

$$\hat{R} = -6 e^{-2f} (2\ddot{f} + 5\dot{f}^2) + e^{-2f} \tilde{R}, \quad (5.25)$$

is divergent, indicating a breakdown of the supergravity approximation in this region. When transformed back to the original variables y and $h(y)$, the solution (5.24) is valid

⁴This expression is obtained from (5.21) by applying the transformation (5.7).




Figure 3. Numerical solution in terms of the original functions $h(y)$ and $\phi(y)$ for case 3 ($\psi_1 = 0$, $\psi_2 = 1$). We have adjusted the free constants to $\alpha' = 4$, $y_0 = 0$ and $\phi_0 = 4$ for concreteness. The solution becomes singular at the point $y = y_0 = 0$, which is the physical location of the domain wall.

only in the half-space $y \in [y_0, \infty)$, where $y_0 = y(\tau_{\text{dw}})$ is the physical location of the domain wall in the y coordinate.

As in the previous case, we obtain a solution for all $y \in \mathbb{R}$ by directly solving (5.2)–(5.5). We find

$$y - y_0 = h - \frac{\sqrt{\alpha'}}{2} \arctan\left(\frac{2h}{\sqrt{\alpha'}}\right). \quad (5.26)$$

This case can also be found in [23], section 4.5.3. Again, the graph of $h(y)$ has a kink shape and a zero at the value $y = y_0$, which indicates the location of the domain wall. For $y \rightarrow \pm\infty$, the function $h(y)$ becomes approximately linear $h(y) \approx y$. The solution for $\phi(y)$ is given implicitly by

$$\phi(h) = \log\left(\frac{h^2}{4h^2 + \alpha'}\right) + \phi_0. \quad (5.27)$$

The graph of $\phi(y)$ has a singularity, $\phi \rightarrow -\infty$, at the domain wall location $y = y_0$ and approaches the limiting value $\phi \rightarrow \log(\frac{1}{4}) + \phi_0$ as $y \rightarrow \pm\infty$. The plots of $h(y)$ and $\phi(y)$ are shown in figure 3. We remark that this case can be obtained from case 2 by applying the transformation $\alpha' \rightarrow -\alpha'$ (and using that $\text{artanh}(iz) = i \arctan(z)$). This is due to the fact that cases 2 and 3 are related by interchanging ψ_1 and ψ_2 , which amounts to a sign flip of the order α' term in (5.14). In the same way, case 5 is related to case 4, and case 7 is related to case 6, as can be seen below.

Case 4: $\psi_1 = \text{kink}$, $\psi_2 = 0$. The solution in this case becomes

$$e^{2f} = e^{2(\tau-\tau_0)} + \frac{\alpha'}{16} [1 - \tanh(\tau-\tau_1)]^2, \quad \phi(\tau) = \phi_0 + 2(f-\tau). \quad (5.28)$$

It is a special case of [22], section 5.2 (with $Q_e = 0$, $a = 1$ and taking the decompactification limit in the S^1 direction). As $\tau \rightarrow +\infty$, the function $f(\tau)$ approaches the linear solution




Figure 4. Exemplary solutions for case 4 ($\psi_1 = \text{kink}$, $\psi_2 = 0$) with $\alpha' = 16$ and $\phi_0 = 10$.

$f(\tau) = \tau$. For $\tau \rightarrow -\infty$, the function $f(\tau)$ converges to the constant value $\frac{1}{2} \log(\frac{\alpha'}{4})$. The graph for finite values of τ qualitatively depends on whether $\tau_0 < \tau_1$ or vice versa. This can be seen from figure 4, where $f(\tau)$ and $\phi(\tau)$ are plotted for different values of the integration constants. We note also that the scalar curvature (5.25) remains finite for all $\tau \in \mathbb{R}$.

Case 5: $\psi_1 = 0$, $\psi_2 = \text{kink}$. The general solution (5.14) becomes

$$e^{2f} = e^{2(\tau-\tau_0)} - \frac{\alpha'}{16} [1 - \tanh(\tau-\tau_2)]^2, \quad \phi(\tau) = \phi_0 + 2(f-\tau). \quad (5.29)$$

The right-hand side approaches zero as $\tau \rightarrow \tau_{\text{dw}}$ from above and then becomes negative for sufficiently small values of τ . Hence, $f(\tau)$ derived from expression (5.29) is ill-defined for $\tau < \tau_{\text{dw}}$. The limiting value τ_{dw} is given by

$$e^{\tau_{\text{dw}}} = \frac{e^{\frac{2}{3}(\tau_0+2\tau_2)} X^{\frac{1}{3}} - 2e^{2\tau_2}}{\sqrt{6} e^{\frac{1}{3}(\tau_0+2\tau_2)} X^{\frac{1}{6}}}, \quad \text{where} \quad X = 27\alpha' + 8e^{2(\tau_2-\tau_0)} + \sqrt{27\alpha' (27\alpha' + 16e^{2(\tau_2-\tau_0)})}. \quad (5.30)$$

In addition, the scalar curvature (5.25) diverges as $\tau \rightarrow \tau_{\text{dw}}$ from above. The limiting value τ_{dw} corresponds to the physical location of the domain wall and our solution exists on the half-space $\tau \in (\tau_{\text{dw}}, \infty)$. The graphs of $f(\tau)$ and $\phi(\tau)$ are displayed in figure 5.

Case 6: $\psi_1 = \text{kink}$, $\psi_2 = 1$. For this set-up, we obtain from (5.14)

$$e^{2f} = e^{2(\tau-\tau_0)} + \frac{\alpha'}{16} [\tanh(\tau-\tau_1) + 1] [\tanh(\tau-\tau_1) - 3]. \quad (5.31)$$

For $\tau \rightarrow +\infty$, this solution behaves approximately linear $f(\tau) \approx \tau$. However, depending on the choice of the free parameters, the function $f(\tau)$ may not be well-defined everywhere, because there can be regions where the right-hand side of (5.31) becomes negative. We have plotted two qualitatively different scenarios in figure 6. In the plot on the left, the function $f(\tau)$ is well-defined everywhere. In addition, the scalar curvature (5.25) has a kink shape implying in particular that it is finite and smooth for all $\tau \in \mathbb{R}$. The plot on the right is very similar to case 5. In this case, the function $f(\tau)$ derived from (5.31) is ill-defined for $\tau < \tau_{\text{dw}}$, where

$$e^{\tau_{\text{dw}}} = \frac{\sqrt{\alpha' e^{2\tau_0} + e^{\tau_0+2\tau_1}} \sqrt{\alpha' e^{\tau_0-3\tau_1}} \sqrt{\alpha' e^{\tau_0-\tau_1} + 16e^{\tau_1-\tau_0}} - 8e^{2\tau_1}}{2\sqrt{2}}, \quad (5.32)$$




Figure 5. Exemplary solution for case 5 ($\psi_1 = 0$, $\psi_2 = \text{kink}$) with $\alpha' = 16$, $\tau_0 = -5$, $\tau_2 = 3$ and $\phi_0 = -2$.




Figure 6. Plots of the solution for case 6 ($\psi_1 = \text{kink}$, $\psi_2 = 1$) for $\alpha' = 44050$ on the left and $\alpha' = 4 \cdot 10^5$ on the right. The other parameters have been adjusted to $\tau_0 = 5$, $\tau_1 = 10$ and $\phi_0 = 0$. The two graphs have the same asymptotics as $\tau \rightarrow +\infty$, but behave qualitatively differently elsewhere.

and the scalar curvature (5.25) diverges as $\tau \rightarrow \tau_{\text{dw}}$ from above. As in case 5, the limiting value τ_{dw} corresponds to the physical location of the domain wall and our solution exists on the half-space $\tau \in (\tau_{\text{dw}}, \infty)$. The distinction between the two scenarios can be made by means of the radicand in the numerator on the right-hand side of expression (5.32). If it is positive, the solution behaves as shown in the plot on the right. This is the case when $\alpha' > 2e^{2(\tau_1 - \tau_0)}$. For $\alpha' \leq 2e^{2(\tau_1 - \tau_0)}$ on the other hand, the solution is globally well-defined as shown in the plot on the left.

Case 7: $\psi_1 = 1$, $\psi_2 = \text{kink}$. The general solution (5.14) now reads as follows,

$$e^{2f} = e^{2(\tau - \tau_0)} - \frac{\alpha'}{16} [\tanh(\tau - \tau_2) + 1] [\tanh(\tau - \tau_2) - 3] . \quad (5.33)$$

This case is similar to case 6, except for the flipped sign in front of the order α' term on the right-hand side. It has already appeared in [1] and in [22], section 5.7 as a special case




Figure 7. Exemplary solution for case 7 ($\psi_1 = 1$, $\psi_2 = \text{kink}$) with $\alpha' = 64$, $\tau_0 = 25$, $\tau_2 = 10$ and $\phi_0 = 20$.

(with $Q_e = 0$, $a = 1$ and taking the decompactification limit in the S^1 direction). The kink solution (3.13) for ψ_2 approaches the value one, as $\tau \rightarrow -\infty$. In this region, the α' corrections cancel and we recover the zeroth order behavior, that is linear f and constant dilaton. In the limit $\tau \rightarrow +\infty$, the expression $(\psi_1^2 - \psi_2^2)$ becomes 1. Thus, for $\tau \gg 1$, the functions f and ϕ also become linear and constant, respectively. However, they are shifted by an offset compared to the $\tau \rightarrow -\infty$ asymptotics. The α' corrections are non-constant in an intermediate region and have the important effect of gluing together the different asymptotic functions. This can be seen from figure 7, where the graphs of $f(\tau)$ and $\phi(\tau)$ are plotted for this case. The scalar curvature (5.25) is finite and smooth for all $\tau \in \mathbb{R}$.

Case 8: $\psi_1 = \text{kink}$, $\psi_2 = \text{kink}$, with $\tau_1 \neq \tau_2$. The general solution (5.14) turns into the following expression,

$$e^{2f} = e^{2(\tau-\tau_0)} + \frac{\alpha'}{16} [\tanh^2(\tau-\tau_1) - 2 \tanh(\tau-\tau_1) - \tanh^2(\tau-\tau_2) + 2 \tanh(\tau-\tau_2)] . \quad (5.34)$$

For $\tau \rightarrow +\infty$, the contributions from the α' corrections vanish and we recover the zeroth order behavior, that is linear f and constant dilaton. For τ finite, the corrections become important and the precise behavior depends on the choice of free parameters. Two qualitatively different scenarios are depicted in figure 8. The plot on the left is an example from a region in parameter space, where the solution resembles that of case 7, with an additional bump, however. In this case the solution is globally well-defined and the scalar curvature (5.25) is finite and smooth everywhere. In other regions of parameter space, the solution is similar to that of case 5. This can be seen from the plot on the right. Here, the function $f(\tau)$ is ill-defined for $\tau < \tau_{\text{dw}}$, where τ_{dw} is the log of the largest real root of the octic equation

$$4(b^2 + x^2)^2 (c^2 + x^2)^2 + \alpha' a^2 (b^2 - c^2) ((b^2 + c^2)x^2 + 2b^2 c^2) = 0 , \quad (5.35)$$




Figure 8. Plots of the solution for case 8 ($\psi_1 = \text{kink}$, $\psi_2 = \text{kink}$, with $\tau_1 \neq \tau_2$) for $\tau_1 = 3$ on the left and $\tau_1 = -6$ on the right. The other parameters have been adjusted to $\alpha' = 64$, $\tau_0 = 7$, $\tau_2 = -5$ and $\phi_0 = -4$. The two graphs have the same asymptotics as $\tau \rightarrow +\infty$, but behave qualitatively differently elsewhere.

with $x = e^{\tau_{\text{dw}}}$, $a = e^{\tau_0}$, $b = e^{\tau_1}$, $c = e^{\tau_2}$. The closed-form expression for τ_{dw} is very lengthy and is thus omitted here for the sake of brevity. The limiting value τ_{dw} corresponds to the physical location of the domain wall and the scalar curvature (5.25) diverges as $\tau \rightarrow \tau_{\text{dw}}$ from above. Our solution exists on the half-space $\tau \in (\tau_{\text{dw}}, \infty)$. On the other hand, if there is no solution of (5.35) over the positive reals, we are in a region of parameter space where a globally well-defined solution, such as the one shown in the plot on the left, exists.

6 Conclusions and outlook

In this paper, we have studied (1+3)-dimensional domain wall solutions of heterotic supergravity on a six-dimensional warped nearly Kähler manifold X_6 in the presence of gravitational and gauge instantons of the type constructed in [1]. The instanton contributions are necessary for solving the Yang-Mills sector and the Bianchi identity (3.18) at order α' , which is the order we have considered. The ten-dimensional solutions constructed in this paper are of the form

$$\hat{g} = \eta_{\alpha\beta} dx^\alpha dx^\beta + dy^2 + (h(y))^2 \tilde{g}_{uv}(x^w) dx^u dx^v, \quad (6.1)$$

$$\hat{H} = H + dy \wedge H_y, \quad (6.2)$$

$$\hat{\phi} = \phi(y). \quad (6.3)$$

where all fields only depend on the non-compact coordinate y transverse to the domain wall. Our solutions preserve two real supercharges, which corresponds to $\mathcal{N}=1/2$ supersymmetry from the viewpoint of the four non-compact dimensions spanned by $\{x^\alpha, y\}$.

Following the general formalism developed in [24, 25], we introduced a pair of y -dependent $\text{SU}(3)$ structure forms (J, Ω) on X_6 defined via the parallel spinors η_\pm and rewrote the BPS equations (2.1) as a set of compatibility relations involving J , Ω , H , H_y and ϕ . There is also a static, that is y -independent, $\text{SU}(3)$ structure denoted $(\tilde{J}, \tilde{\Omega})$ on X_6 . The two structures (J, Ω) and $(\tilde{J}, \tilde{\Omega})$ are related by means of the warp factor $h(y)$ and a y -dependent mixing angle β .

The BPS equations and Bianchi identity then reduce to a set of ordinary differential equations involving the free functions h , β and ϕ . The complete system of coupled non-linear ordinary differential equations, as summarized in (4.21)–(4.26), is too complicated to solve in full generality. Instead, we studied the special branches $W_1^+ = 0$ and $W_1^- = 0$. While the case $W_1^+ = 0$ can be quickly discarded for it leads to a singular metric, the second case, $W_1^- = 0$, allows for the construction of a variety of solutions depending on the choice of instantons $\psi_{1,2}$ for the gravitational and gauge sector. In total, there are eight distinct cases, including already known solutions (cases 1-4 and 7) from [1, 22, 23] as well as some new ones (cases 5, 6 and 8).

For the solutions with $W_1^- = 0$, the ten-dimensional fields take the following simpler form,

$$\hat{g} = \eta_{\alpha\beta} dx^\alpha dx^\beta + e^{2f(\tau)} (d\tau^2 + \tilde{g}_{uv}(x^w) dx^u dx^v) , \quad (6.4)$$

$$\hat{H} = -\frac{\alpha'}{4} (\psi_1^2(2\psi_1 - 3) - \psi_2^2(2\psi_2 - 3)) \tilde{\Omega}_+ , \quad (6.5)$$

$$\phi = \phi_0 + 2(f - \tau) , \quad (6.6)$$

$$e^{2f} = e^{2(\tau - \tau_0)} + \frac{\alpha'}{4} (\psi_1^2 - \psi_2^2) , \quad (6.7)$$

where $\psi_{1,2}$ are either 0, 1 or the kink solution (3.13), and we used the convenient reparameterization $dy = e^{f(\tau)} d\tau$, $h(y) = e^{f(\tau(y))}$. In case one (equivalent to zeroth order in α'), $f = \tau - \tau_0$, which leads to a cone metric with constant dilaton and vanishing NS 3-form flux. The other cases typically asymptote to this zeroth order behavior either at $\tau \rightarrow +\infty$, $\tau \rightarrow -\infty$ or $\tau \rightarrow \pm\infty$. Close to the domain wall, care must be taken as to the validity of the supergravity approximation. Indeed, in some of the cases, the scalar curvature diverges as the domain wall is approached.

The NS 3-form flux \hat{H} is always proportional to $\tilde{\Omega}_+$ in all our explicit constructions. It would be interesting to have access to solutions with a more general \hat{H} that also includes terms proportional to $\tilde{\Omega}_-$ and $\tilde{J} \wedge dy$. It remains to be seen whether this can be achieved by finding a solution of the general system (4.21)–(4.26) with both $W_1^+ \neq 0$ and $W_1^- \neq 0$ or whether another, perhaps rather different, ansatz is necessary. This is left for future work.

Acknowledgments

We would like to thank Alexander Popov for useful discussions and for collaboration in the early stages of this work. We are grateful to the referee for careful reading and constructive comments. This work was partially supported by the Deutsche Forschungsgemeinschaft grant LE 838/13. A.S.H. thanks the Albert-Einstein-Institute and its director Hermann Nicolai for warm hospitality and generous financial support through the award of a Max Planck Society research stipend. The work of E.T.M. was supported by the Riemann Center for Geometry and Physics, École Normale Supérieure de Lyon and by the Albert-Einstein-Institute. E.T.M. personally thanks Henning Samtleben and Hermann Nicolai for warm hospitality during completion of part of this project.

A Conventions

In this paper we use the following conventions for indices and normalizations. The full range of (1+9)-dimensional indices is split by the presence of the domain wall and will be distinguished by means of the following set of Greek and Latin letters,

$$\begin{aligned}\mu, \nu, \rho, \sigma &= 0, 1, \dots, 9, \\ \alpha, \beta, \gamma, \delta, \epsilon &= 0, 1, 2, \\ a, b, c, \dots, m, n &= 3, 4, \dots, 9, \\ u, v, w, x, y, z &= 4, 5, \dots, 9.\end{aligned}\tag{A.1}$$

These are understood as curved space-time indices. In addition, we sometimes need to use tangent space (local Lorentz) indices, which are denoted by underlined indices.

(Anti-)symmetrization is always performed with a factor of $(1/n!)$, that is with weight one. For example, we define

$$A_{[\mu}B_{\nu]} := \frac{1}{2!}(A_{\mu}B_{\nu} - A_{\nu}B_{\mu}),\tag{A.2}$$

for the case $n = 2$. A p -form ω is expanded into components according to

$$\omega := \frac{1}{p!}\omega_{\mu_1\dots\mu_p}dx^{\mu_1} \wedge \dots \wedge dx^{\mu_p}.\tag{A.3}$$

The Clifford action of a p -form ω on a spinor ϵ is defined as

$$\omega \cdot \epsilon := \frac{1}{p!}\omega_{\mu_1\dots\mu_p}\gamma^{\mu_1\dots\mu_p}\epsilon,\tag{A.4}$$

where $\gamma^{\mu_1\dots\mu_p} := \gamma^{[\mu_1} \dots \gamma^{\mu_p]}$ and we use the Clifford algebra convention $\{\gamma^{\mu}, \gamma^{\nu}\} = 2g^{\mu\nu}$ for the higher-dimensional gamma matrices γ^{μ} .

A connection ∇ on a manifold \mathcal{M} with vielbein σ_b^a is defined to act on vectors v^b and spinors ϵ in the following way,

$$\begin{aligned}\nabla_a v^b &= \partial_a v^b + \Gamma_a{}^b{}_c v^c, \\ \nabla_a \epsilon &= \partial_a \epsilon + \frac{1}{4}\omega_a \cdot \epsilon, \quad \omega_a = \frac{1}{2}\omega_a{}^{\underline{c}}{}_{\underline{b}}\gamma_{\underline{c}}{}^{\underline{b}}.\end{aligned}\tag{A.5}$$

The components of the spin connection $\omega_a{}^{\underline{b}}{}_{\underline{c}}$ are related to $\Gamma_a{}^b{}_c$ via

$$\omega_a{}^{\underline{b}}{}_{\underline{c}} = \Gamma_a{}^{\underline{b}}{}_{\underline{c}} - \sigma_{\underline{c}}^d \partial_a \sigma_d^{\underline{b}}.\tag{A.6}$$

The torsion T^a of the connection ∇ is defined as $T^a = \nabla \sigma^a$. Using (A.5)–(A.6), we can expand the torsion into components as follows,

$$\begin{aligned}T^a &= \nabla \sigma^a = d\sigma^a + \omega_a{}^{\underline{b}} \wedge \sigma^{\underline{b}} = \Gamma_{\underline{b}\underline{c}}^a \sigma^{\underline{b}} \wedge \sigma^{\underline{c}}, \quad \text{or} \\ T_{bc}{}^a &= \Gamma_{[b}{}^a{}_{c]},\end{aligned}\tag{A.7}$$

where $\omega_a{}^{\underline{b}} = \omega_c{}^{\underline{a}}{}_{\underline{b}} dx^c$ is the connection 1-form and the $\sigma^a = \sigma_b^a dx^b$ define an orthonormal frame on \mathcal{M} .

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] D. Harland and C. Nolle, *Instantons and Killing spinors*, *JHEP* **03** (2012) 082 [[arXiv:1109.3552](https://arxiv.org/abs/1109.3552)] [[INSPIRE](#)].
- [2] M. Graña, *Flux compactifications in string theory: a comprehensive review*, *Phys. Rept.* **423** (2006) 91 [[hep-th/0509003](https://arxiv.org/abs/hep-th/0509003)] [[INSPIRE](#)].
- [3] B. Wecht, *Lectures on nongeometric flux compactifications*, *Class. Quant. Grav.* **24** (2007) S773 [[arXiv:0708.3984](https://arxiv.org/abs/0708.3984)] [[INSPIRE](#)].
- [4] M.R. Douglas and S. Kachru, *Flux compactification*, *Rev. Mod. Phys.* **79** (2007) 733 [[hep-th/0610102](https://arxiv.org/abs/hep-th/0610102)] [[INSPIRE](#)].
- [5] R. Blumenhagen, B. Körs, D. Lüst and S. Stieberger, *Four-dimensional string compactifications with D-branes, orientifolds and fluxes*, *Phys. Rept.* **445** (2007) 1 [[hep-th/0610327](https://arxiv.org/abs/hep-th/0610327)] [[INSPIRE](#)].
- [6] H. Samtleben, *Lectures on gauged supergravity and flux compactifications*, *Class. Quant. Grav.* **25** (2008) 214002 [[arXiv:0808.4076](https://arxiv.org/abs/0808.4076)] [[INSPIRE](#)].
- [7] F. Quevedo, *Lectures on superstring phenomenology*, *AIP Conf. Proc.* **359** (1996) 202 [[hep-th/9603074](https://arxiv.org/abs/hep-th/9603074)] [[INSPIRE](#)].
- [8] D. Joyce, *Lectures on Calabi-Yau and special Lagrangian geometry*, [math/0108088](https://arxiv.org/abs/math/0108088) [[INSPIRE](#)].
- [9] M.B. Green and J.H. Schwarz, *Anomaly cancellation in supersymmetric $D = 10$ gauge theory and superstring theory*, *Phys. Lett. B* **149** (1984) 117 [[INSPIRE](#)].
- [10] D.J. Gross, J.A. Harvey, E.J. Martinec and R. Rohm, *Heterotic string theory. 1. The free heterotic string*, *Nucl. Phys. B* **256** (1985) 253 [[INSPIRE](#)].
- [11] D.J. Gross, J.A. Harvey, E.J. Martinec and R. Rohm, *Heterotic string theory. 2. The interacting heterotic string*, *Nucl. Phys. B* **267** (1986) 75 [[INSPIRE](#)].
- [12] C.M. Hull, *Anomalies, ambiguities and superstrings*, *Phys. Lett. B* **167** (1986) 51 [[INSPIRE](#)].
- [13] S. Ivanov, *Heterotic supersymmetry, anomaly cancellation and equations of motion*, *Phys. Lett. B* **685** (2010) 190 [[arXiv:0908.2927](https://arxiv.org/abs/0908.2927)] [[INSPIRE](#)].
- [14] E.A. Bergshoeff and M. de Roo, *The quartic effective action of the heterotic string and supersymmetry*, *Nucl. Phys. B* **328** (1989) 439 [[INSPIRE](#)].
- [15] K. Becker and S. Sethi, *Torsional heterotic geometries*, *Nucl. Phys. B* **820** (2009) 1 [[arXiv:0903.3769](https://arxiv.org/abs/0903.3769)] [[INSPIRE](#)].
- [16] G.T. Horowitz and A. Strominger, *Black strings and P-branes*, *Nucl. Phys. B* **360** (1991) 197 [[INSPIRE](#)].
- [17] A. Strominger, *Heterotic solitons*, *Nucl. Phys. B* **343** (1990) 167 [Erratum *ibid.* **B** **353** (1991) 565] [[INSPIRE](#)].
- [18] J.A. Harvey and A. Strominger, *Octonionic superstring solitons*, *Phys. Rev. Lett.* **66** (1991) 549 [[INSPIRE](#)].

- [19] R.R. Khuri, *Remark on string solitons*, *Phys. Rev. D* **48** (1993) 2947 [[hep-th/9305143](#)] [[INSPIRE](#)].
- [20] M. Günaydin and H. Nicolai, *Seven-dimensional octonionic Yang-Mills instanton and its extension to an heterotic string soliton*, *Phys. Lett. B* **351** (1995) 169 [Addendum *ibid.* **B 376** (1996) 329] [[hep-th/9502009](#)] [[INSPIRE](#)].
- [21] E.K. Loginov, *Some comments on string solitons*, *Phys. Rev. D* **77** (2008) 105003 [[arXiv:0805.0870](#)] [[INSPIRE](#)].
- [22] K.-P. Gemmer, A.S. Haupt, O. Lechtenfeld, C. Nölle and A.D. Popov, *Heterotic string plus five-brane systems with asymptotic AdS_3* , *Adv. Theor. Math. Phys.* **17** (2013) 771 [[arXiv:1202.5046](#)] [[INSPIRE](#)].
- [23] M. Klaput, A. Lukas, C. Matti and E.E. Svanes, *Moduli stabilising in heterotic nearly Kähler compactifications*, *JHEP* **01** (2013) 015 [[arXiv:1210.5933](#)] [[INSPIRE](#)].
- [24] J. Gray, M. Larfors and D. Lüst, *Heterotic domain wall solutions and $SU(3)$ structure manifolds*, *JHEP* **08** (2012) 099 [[arXiv:1205.6208](#)] [[INSPIRE](#)].
- [25] A. Lukas and C. Matti, *G-structures and domain walls in heterotic theories*, *JHEP* **01** (2011) 151 [[arXiv:1005.5302](#)] [[INSPIRE](#)].
- [26] S. Chiossi and S. Salamon, *The intrinsic torsion of $SU(3)$ and G_2 structures*, [math/0202282](#) [[INSPIRE](#)].
- [27] N.J. Hitchin, *Stable forms and special metrics*, [math/0107101](#) [[INSPIRE](#)].
- [28] C. Mayer and T. Mohaupt, *Domain walls, Hitchin's flow equations and G_2 -manifolds*, *Class. Quant. Grav.* **22** (2005) 379 [[hep-th/0407198](#)] [[INSPIRE](#)].
- [29] J. Louis and S. Vaula, *$N = 1$ domain wall solutions of massive type-II supergravity as generalized geometries*, *JHEP* **08** (2006) 058 [[hep-th/0605063](#)] [[INSPIRE](#)].
- [30] P. Smyth and S. Vaula, *Domain wall flow equations and $SU(3) \times SU(3)$ structure compactifications*, *Nucl. Phys. B* **828** (2010) 102 [[arXiv:0905.1334](#)] [[INSPIRE](#)].
- [31] J.P. Gauntlett, D. Martelli, S. Pakis and D. Waldram, *G structures and wrapped NS5-branes*, *Commun. Math. Phys.* **247** (2004) 421 [[hep-th/0205050](#)] [[INSPIRE](#)].
- [32] D. Martelli and J. Sparks, *Non-Kähler heterotic rotations*, *Adv. Theor. Math. Phys.* **15** (2011) 131 [[arXiv:1010.4031](#)] [[INSPIRE](#)].
- [33] X. de la Ossa and E.E. Svanes, *Holomorphic bundles and the moduli space of $N = 1$ supersymmetric heterotic compactifications*, *JHEP* **10** (2014) 123 [[arXiv:1402.1725](#)] [[INSPIRE](#)].
- [34] M. Fernandez, S. Ivanov, L. Ugarte and D. Vassilev, *Non-Kähler heterotic string solutions with non-zero fluxes and non-constant dilaton*, *JHEP* **06** (2014) 073 [[arXiv:1402.6107](#)] [[INSPIRE](#)].
- [35] I.V. Melnikov, R. Minasian and S. Sethi, *Heterotic fluxes and supersymmetry*, *JHEP* **06** (2014) 174 [[arXiv:1403.4298](#)] [[INSPIRE](#)].
- [36] T. Maxfield and S. Sethi, *Domain walls, triples and acceleration*, *JHEP* **08** (2014) 066 [[arXiv:1404.2564](#)] [[INSPIRE](#)].
- [37] A. Gray, *The structure of nearly Kähler manifolds*, *Math. Ann.* **223** (1976) 233.